arXiv:1209.2471v1 [math.CO] 12 Sep 2012

Every 4-regular graph is acyclically edge-6-colorable

Weifan Wang*, Qiaojun Shu
Department of Mathematics, Zhejiang Normal University, Jinhua 321004, China

Yiqiao Wang
School of Management, Beijing University of Chinese Medicine, Beijing 100029, China

September 13, 2012

Abstract

An acyclic edge coloring of a graph G is a proper edge coloring such that no bichromatic cycles are produced. The acyclic chromatic index $a'(G)$ of G is the smallest integer k such that G has an acyclic edge coloring using k colors. Fiamčik (1978) and later Alon, Sudakov and Zaks (2001) conjectured that $a'(G) \leq \Delta + 2$ for any simple graph G with maximum degree Δ. Basavaraju and Chandran (2009) showed that every graph G with $\Delta = 4$, which is not 4-regular, satisfies the conjecture. In this paper, we settle the 4-regular case, i.e., we show that every 4-regular graph G has $a'(G) \leq 6$.

Keywords: acyclic edge coloring; 4-regular graph; maximum degree

AMS subject classification. 05C15

1 Introduction

Only simple graphs are considered in this paper. Let G be a graph with vertex set $V(G)$ and edge set $E(G)$. A *proper edge-k-coloring* is a mapping $c : E(G) \to \{1, 2, \ldots, k\}$ such that any two adjacent edges receive different colors. The graph G is *edge-k-colorable* if it has an edge-k-coloring. The *chromatic index* $\chi'(G)$ of G is the smallest integer k such that G is edge-k-colorable. A proper edge-k-coloring of G is called *acyclic* if there are no bichromatic cycles in G, i.e., the union of any two color classes induces a subgraph of G that is a forest. The *acyclic chromatic index* of G, denoted $a'(G)$, is the smallest integer k such that G is acyclically edge-k-colorable.

---

*Research supported partially by NSFC(No.11071223) and ZJNSF(No.Z6090150); Corresponding author. Email: wwf@zjnu.cn.

Let $\Delta(G)$ (Δ for short) denote the maximum degree of a graph G. By Vizing's Theorem [26], $\Delta \leq \chi'(G) \leq \Delta + 1$. Thus, it holds trivially that $a'(G) \geq \chi'(G) \geq \Delta$. Fiamčik [10] and later Alon, Sudakov and Zaks [1] made the following conjecture:

Conjecture 1 *For any graph G, $a'(G) \leq \Delta + 2$.*

Using probabilistic method, Alon, McDiarmid and Reed [2] proved that $a'(G) \leq 64\Delta$ for any graph G. This bound has been recently improved to that $a'(G) \leq 16\Delta$ in [15], that $a'(G) \leq \lceil 9.62(\Delta - 1)\rceil$ in [16], and that $a'(G) \leq 4\Delta$ in [9]. In [17], Něsetřil and Wormald proved that $a'(G) \leq \Delta + 1$ for a random Δ-regular graph G. The acyclic edge coloring of some special classes of graphs was also investigated, including subcubic graphs [3, 18], outerplanar graphs [13, 14], series-parallel graphs [12, 22], 2-degenerate graphs [5], and planar graphs [6, 7, 8, 11, 19, 23, 25, 27]. In particular, Conjecture 1 was confirmed for planar graphs without 3-cycles [20], or without 4-cycles [24], or without 5-cycles [21]. It was shown in [6] that every planar graph G has $a'(G) \leq \Delta + 12$. This upper bound was recently improved to that $a'(G) \leq \Delta + 7$ [25].

Basavaraju and Chandran [4] showed that if G is a graph with $\Delta = 4$ and $|E(G)| \leq 2|V(G)| - 1$, then $a'(G) \leq 6$. Equivalently, every non-regular graph G with $\Delta = 4$ satisfies Conjecture 1. In this paper, we will settle the case where G is 4-regular. Our result and the result of [4] confirm Conjecture 1 for graphs with $\Delta = 4$.

2 Acyclic chromatic indices

In this section, we discuss the acyclic chromatic indices of 4-regular graphs. Before establishing our main result, we need to introduce some notation.

Assume that c is a partial acyclic edge-k-coloring of a graph G using the color set $C = \{1, 2, \ldots, k\}$. For a vertex $v \in V(G)$, we use $C(v)$ to denote the set of colors assigned to edges incident to v under c. If the edges of a cycle $ux \ldots vu$ are alternatively colored with colors i and j, then we call such cycle an $(i,j)_{(u,v)}$-*cycle*. If the edges of a path $ux \ldots v$ are alternatively colored with colors i and j, then we call such path an $(i,j)_{(u,v)}$-*path*. In the following Figures, black spots are pairwise distinct, whereas the others may be not.

For simplicity, we use $\{e_1, e_2, \cdots, e_m\} \to a$ to express that all edges e_1, e_2, $\cdots$, e_m are colored or recolored with the color a. In particular, when $m = 1$, we write simply $e_1 \to a$. Moreover, we use $(e_1, e_2, \cdots, e_m)_c = (a_1, a_2, \cdots, a_m)$ to denote the fact that

$c(e_i) = a_i$ for $i = 1, 2, \dots, m$. Let $(e_1, e_2, \cdots, e_n) \to (b_1, b_2, \cdots, b_n)$ denote the fact that e_i is colored or recolored with the color b_i for $i = 1, 2, \dots, n$.

Now we give two useful lemmas, which will be used in the following discussion.

Lemma 1 *Suppose that a graph G has an edge-6-coloring c such that $c(uv) = i$, where uv is an arbitrary edge of G. If uv can be recolored properly with the color j $(\neq i)$, then G contains no $(i, j)_{(u,v)}$-path under c.*

Lemma 1 holds obviously by the definition.

Lemma 2 *Suppose that a graph G has an edge-6-coloring c. Let $P = uv_1v_2 \dots v_kv_{k+1}$ be an $(i, j)_{(u,v_{k+1})}$-path in G with $c(uv_1) = i$ and $j \notin C(u)$. If $w \notin V(P)$, then there does not exist an $(i, j)_{(u,w)}$-path in G under c.*

Proof. Otherwise, assume that there is an $(i, j)_{(u,w)}$-path $Q = uw_1w_2 \dots w_m$ in G, where $w = w_m$ and $m \geq 1$. Note that some w_s may be identical to some v_t, $1 \leq s \leq m-1$, $1 \leq t \leq k+1$. Since $j \notin C(u)$ and $w \notin V(P)$, it is easy to see that there exists a vertex v_t and some vertex w_s such that v_tv_{t+1} and v_tw_s have the same color i or j, which contradicts the fact that c is a proper edge coloring. $\square$

Theorem 1 *If G is a 4-regular graph without 3-cycles, then $a'(G) \leq 6$.*

Proof. By Vizing's theorem [26], G admits a proper edge-6-coloring c using the color set $C = \{1, 2, \dots, 6\}$. Let $\tau(c)$ denote the number of bichromatic cycles in G with respect to the coloring c. If $\tau(c) = 0$, then c is an acyclic edge-6-coloring of G, hence we are done. Otherwise, $\tau(c) > 0$. Let B be a bichromaic cycle of G. We are going to show that there is a proper edge-6-coloring c', formed by recoloring suitably some edges of G, such that B is no longer a bichromatic cycle and on new bichromatic cycles are produced. Namely, $\tau(c') < \tau(c)$. Repeating this process, we finally obtain an acyclic edge-6-coloring c^* of G.

To arrive at our conclusion, assume, w.l.o.g., that B is a $(4, 1)_{(u,v)}$-cycle with $c(uv) = 1$. Let u_1, u_2, u_3 be the neighbors of u different from v, and v_1, v_2, v_3 the neighbors of v different from u. Since G contains no triangles, $u_1, u_2, u_3, v_1, v_2, v_3$ are pairwise distinct. Clearly, we may assume that $c(uu_2) = c(vv_2) = 4$. Since $2 \leq |C(u) \cap C(v)| \leq 4$, the proof is split into the following three cases, as shown in Fig. 1.

Case 1 $|C(u) \cap C(v)| = 2$, say $(uu_1, uu_3)_c = (2, 5)$ and $(vv_1, vv_3)_c = (3, 6)$.

By symmetry, we may assume that $C(u_2) \in \{\{1, 4, 3, 6\}, \{1, 4, 2, 6\}, \{1, 4, 2, 5\}\}$.



$|C(u)\cap C(v)|=2$ $|C(u)\cap C(v)|=3$ $|C(u)\cap C(v)|=4$

Fig. 1: Three cases on $|C(u)\cap C(v)|=2,3,4$.

Case 1.1 $C(u_2)=\{1,4,3,6\}$.

Assume that $3\notin C(u_1)\cup C(u_3)$. If G contains no $(4,3)_{(u_1,u_2)}$-path, then let $(uu_1,uv)\to(3,2)$. Otherwise, G contains a $(4,3)_{(u_1,u_2)}$-path and let $(uu_3,uv)\to(3,5)$. Otherwise, assume that $3\in C(u_1)\cup C(u_3)$ and $6\in C(u_1)\cup C(u_3)$. If $1\notin C(u_1)$ and G contains no $(1,5)_{(u_1,u_3)}$-path, then let $(uu_1,uv)\to(1,2)$. If $4\notin C(u_1)$ and G contains no $(4,5)_{(u_1,u_3)}$-path, then let $(uu_1,uu_2)\to(4,2)$. By Lemma 2, no new bichromatic cycles are produced. Otherwise, assume that:

$(*_{\mathbf{1.1}})$

- (a) $\{1,4\}\subseteq C(u_1)$, *or* G *contains a* $(5,i)_{(u_1,u_3)}$*-path for any* $i\in\{1,4\}\backslash C(u_1)$*;*
- (b) $\{1,4\}\subseteq C(u_3)$, *or* G *contains a* $(2,i)_{(u_1,u_3)}$*-path for any* $i\in\{1,4\}\backslash C(u_3)$*;*
- (c) $\{1,4,3,6\}\subseteq C(u_1)\cup C(u_3)$*;*
- (d) $5\in C(u_1)$ *if* $\{1,4\}\backslash C(u_1)\neq\emptyset$ *and* $2\in C(u_3)$ *if* $\{1,4\}\backslash C(u_3)\neq\emptyset$.

(1.1.1) Assume that $2\notin C(u_3)$.

By $(*_{\mathbf{1.1}})$, $\{1,4\}\subseteq C(u_3)$. If $1\notin C(u_1)$, then let $(uu_1,uu_3,uv)\to(1,2,5)$. If $4\notin C(u_1)$, then let $(uu_1,uu_2,uu_3)\to(4,5,2)$. Thus, $\{1,4\}\subseteq C(u_1)$ and assume that $C(u_1)=\{1,2,3,4\}$ and $C(u_3)=\{5,4,1,6\}$ since $\{3,6\}\subseteq C(u_1)\cup C(u_3)$ by $(*_{\mathbf{1.1}})$. If $1\notin C(v_1)$, then let $(uu_2,uu_3)\to(5,3)$. If $1\notin C(v_3)$, then let $(uu_1,uu_2)\to(6,2)$. Otherwise, $1\in C(v_1)\cap C(v_3)$.

Suppose that we can recolor some edges in $\{vv_1,vv_2,vv_3\}$ such that $C(u)\cap C(v)\in\{\{1,2\},\{1,5\}\}$, or $C(u)\cap C(v)\in\{\{1,2,4\},\{1,5,4\}\}$ and $c(vv_2)\neq 4$, and no new bichromatic cycles are produced in $G-uv$. By Lemma 2, B does not exist even if $4\in C(v)$. If G contains a $(1,i)_{(u,v)}$-cycle for some $i\in\{2,5\}$, then let $(uu_1,uu_3)\to(5,2)$ and no new bichromatic cycles are produced. Next, we claim that:

$(\text{※}_{\mathbf{1}})$ *If we can recolor some edges in* $\{vv_1,vv_2,vv_3\}$ *such that* $C(u)\cap C(v)\in\{\{1,2\},\{1,5\}\}$, *or* $C(u) \cap C(v) \in \{\{1,2,4\},\{1,5,4\}\}$ *and* $c(vv_2) \neq 4$*, and no new bichromatic cycles are produced except the* $(1,i)_{(u,v)}$*-cycle for some* $i \in \{2,5\}$*, we are done.*

- Assume that $C(v_2) = \{4,1,2,5\}$.

If G contains no $(3,i)_{(v_1,v_3)}$-path for some $i \in \{2,5,4\} \backslash C(v_3)$, then let $(vv_3, vv_2) \to (i,6)$ by Lemma 2. Otherwise, G contains a $(3,i)_{(v_1,v_3)}$-path for every $i \in \{2,5,4\} \backslash C(v_3)$ and a $(6,i)_{(v_1,v_3)}$-path for every $i \in \{2,5,4\} \backslash C(v_1)$ similarly. Thus, $3 \in C(v_3)$ and $6 \in C(v_1)$. However, $\{2,5,4\} \subseteq C(v_1) \cup C(v_3)$, a contradiction.

- Assume that $\{2,5\} \backslash C(v_2) \neq \emptyset$ and $2 \notin C(v_2)$. (If $5 \notin C(v_2)$, we can first let $(uu_1, uu_3)_c = (5,2)$ and have a similar discussion.)

By symmetry, $C(v_2) = \{4,1,3,5\}$ or $C(v_2) = \{4,1,3,6\}$.

(i) $C(v_2) = \{4,1,3,6\}$.

If G contains neither a $(2,3)_{(v_1,v_2)}$-path nor a $(2,6)_{(v_2,v_3)}$-path, then let $vv_2 \to 2$ by $(\divideontimes_{\mathbf{1}})$. Otherwise, G contains a $(2,3)_{(v_1,v_2)}$-path and $2 \in C(v_1)$. Assume that $2 \notin C(v_3)$. Then first let $vv_3 \to 2$. Next, if G contains no $(4,6)_{(u_2,v)}$-path, then let $uv \to 6$; otherwise, G contains a $(4,6)_{(u_2,v)}$-path, we let $(uu_1, uu_3, uv) \to (6,2,5)$. Assume that $2 \in C(v_3)$ and $5 \in C(v_1) \cup C(v_3)$ similarly (It follows that $\{2,5\} \subseteq C(v_1) \cup C(v_3)$ if G contains a $(2,6)_{(v_2,v_3)}$-path). Then, let $(vv_1, vv_2, vv_3) \to (4,2,3)$ by $(\divideontimes_{\mathbf{1}})$.

(ii) $C(v_2) = \{4,1,3,5\}$.

If G contains no $(2,3)_{(v_1,v_2)}$-path, then let $vv_2 \to 2$. Otherwise, G contains a $(2,3)_{(v_1,v_2)}$-path and $2 \in C(v_1)$. If $2 \notin C(v_3)$, then let $(vv_3, uv) \to (2,6)$. Otherwise, $2 \in C(v_3)$. If $4 \notin C(v_1)$ and G contains no $(4,6)_{(v_1,v_3)}$-path, then let $(vv_1, vv_2, uv) \to (4,2,3)$. Otherwise, $4 \in C(v_1)$ or G contains a $(4,6)_{(v_1,v_3)}$-path. If $4 \notin C(v_1)$ and G contains a $(4,6)_{(v_1,v_3)}$-path, i.e., $C(v_1) = \{3,1,2,6\}$, $C(v_3) = \{6,1,2,4\}$, then let $(vv_1, vv_2, vv_3, uv) \to (4,6,5,3)$. Otherwise, $C(v_1) = \{3,1,2,4\}$. If $5 \notin C(v_3)$, then let $(vv_2, vv_3) \to (6,5)$; otherwise, $C(v_3) = \{6,1,2,5\}$, let $(vv_2, vv_3) \to (6,4)$ and we are done by $(\divideontimes_{\mathbf{1}})$ or Lemma 2.

(1.1.2) Assume that $2 \in C(u_3)$ and $5 \in C(u_1)$.

Since $\{1,4,3,6\} \subseteq C(u_1) \cup C(u_3)$ by $(*_{\mathbf{1.1}})$, we conclude that $C(u_1) = \{2,5,1,3,5\}$ and $C(u_3) = \{5,2,4,6\}$, or $C(u_1) = \{2,5,1,4\}$ and $C(u_3) = \{5,2,3,6\}$. Let $x_1, x_2, x_3 \neq u$ be the other three neighbors of u_1 and $c(u_1x_1) = 1$. Note that G contains a $(1,2)_{(x_1,u_3)}$-path and $2 \in C(x_1)$ by $(*_{\mathbf{1.1}})$.

- Assume that $C(u_1) = \{2,5,1,3\}$, $C(u_3) = \{5,2,4,6\}$, and $c(u_1x_2) = 3$

If G contains no $(5,6)_{(u_1,u_3)}$ -path, then let $(uu_1, uv) \to (6,2)$. Otherwise, G contains a $(5,6)_{(u_1,u_3)}$-path. If G contains no $(3,2)_{(x_2,u_3)}$-path, then let $(uu_2, uu_3) \to (5,3)$; otherwise, assume that G contains a $(3,2)_{(x_2,u_3)}$ and $2 \in C(x_2)$. If $\{4,6\} \backslash C(x_i) \neq \emptyset$ for some $i \in \{1,2\}$, then let $(uu_2, uu_3, uv) \to (5, c(u_1x_i), 2)$ and recolor uu_1 with a color in $\{4,6\}\backslash C(x_i)$. Otherwise, $C(x_1)=\{1,2,4,6\}$ and $C(x_2)=\{3,2,4,6\}$. Then, we switch the colors of u_1x_1 and u_1x_2, and let $(uu_3,uv)\rightarrow(1,5)$.

- Assume that $C(u_1)=\{2,1,5,4\}$, $C(u_3)=\{5,2,3,6\}$, and $c(u_1x_2)=4$.

Note that G contains a $(4,2)_{(u_1,u_3)}$-path by $(*_{\mathbf{1.1}})$ and then $2\in C(x_2)$. Assume that $3\notin C(x_2)$. If G contains no $(3,5)_{(u_1,u_3)}$-path, then let $(uu_1,uv)\rightarrow(3,2)$; otherwise, G contains a $(3,5)_{(u_1,u_3)}$-path and then let $(uu_1,uu_2,uu_3,uv)\rightarrow(3,5,4,2)$. Otherwise, $3\in C(x_2)$ and $6\in C(x_2)$ similarly. Thus, $C(x_2)=\{4,2,3,6\}$. Assume that $3\notin C(x_1)$ and first let $uu_1\rightarrow 3$. Next, if G contains no $(3,4)_{(u_1,u_2)}$-path, then let $(uu_3,uv)\rightarrow(1,2)$; otherwise, G contains a $(3,4)_{(u_1,u_2)}$-path, we let $(uu_2,uu_3,uv)\rightarrow(2,4,1)$. Otherwise, $3\in C(x_1)$ and $6\in C(x_1)$ similarly. Thus, $C(x_1)=\{1,2,3,6\}$. Then, switch the colors of u_1x_1 and u_1x_2 and let $(uu_3,uv)\rightarrow(1,5)$.

By $(*_{\mathbf{1.1}})$ and Lemma 2, no new bichromatic cycles are produced, as shown in Fig. 2.



Fig. 2: The coloring in Case 1.1.2.

Case 1.2 $C(u_2)=\{1,4,2,6\}$.

If $C(v_2)=\{1,4,2,5\}$, we reduce the proof to Case 1.1. Thus, $\{2,5\}\backslash C(v_2)\neq\emptyset$. If G contains no $(i,2)_{(u_1,u_3)}$-path for some $i\in\{1,3\}\backslash C(u_3)$, then let $(uu_3,uv)\rightarrow(i,5)$. Otherwise, assume that:

$(*_{\mathbf{1.2}})$

(a) $\{1,3\}\subseteq C(u_3)$, *or* G *contains a* $(i,2)_{(u_1,u_3)}$*-path for any* $i\in\{1,3\}\backslash C(u_3)$;

(b) $\{1,3\}\subseteq C(u_3)\cup C(u_1)$, *and* $2\in C(u_3)$ *if* $\{1,3\}\backslash C(u_3)\neq\emptyset$.

Case 1.2.1 $2\notin C(u_3)$.

Then $\{1,3\}\subseteq C(u_3)$ by $(*_{\mathbf{1.2}})$. If $4\notin C(u_1)$, then let $(uu_1,uu_2,uu_3)\rightarrow(4,5,2)$. If $1\notin C(u_1)$ and G contains no $(4,2)_{(u_2,u_3)}$-path, then let $(uu_1,uu_3,uv)\rightarrow(1,2,5)$. Otherwise, $1\in C(u_1)$ or G contains a $(4,2)_{(u_2,u_3)}$-path.

- Assume that $C(u_3) = \{5, 1, 3, 6\}$.

Clearly, $1 \in C(u_1)$ and first let $uu_3 \to 4$. Next, if $5 \notin C(u_1)$, then let $uu_2 \to 5$ and we are done by Lemma 2. Otherwise, $C(u_1) = \{2, 1, 4, 5\}$, we let $(uu_2, uv) \to (3, 5)$.

- Assume that $C(u_3) = \{5, 1, 3, 4\}$.

If G contains no $(4, 6)_{(u_2,u_3)}$-path, then let $(uu_3, uv) \to (6, 5)$. Otherwise, assume that G contains a $(4, 6)_{(u_2,u_3)}$-path. We have to handle two possibilities:

(i) $1 \notin C(u_1)$.

It follows that G contains a $(4, 2)_{(u_2,u_3)}$-path. If $5 \notin C(u_1)$, then let $(uu_1, uv) \to (1, 2)$. If $6 \notin C(u_1)$, then let $(uu_1, uv) \to (6, 2)$. Otherwise, $C(u_1) = \{2, 4, 5, 6\}$ and first let $uu_3 \to 6$. If G contains no $(3, 6)_{(u_1,u_3)}$-path, let $(uu_1, uu_2, uv) \to (3, 5, 2)$; otherwise, let $(uu_2, uv) \to (3, 5)$.

(ii) $1 \in C(u_1)$.

First assume that $3, 5 \notin C(u_1)$. Let $uu_1 \to 3$. If G contains no $(2, 4)_{(u_2,v_2)}$-path, we let $uv \to 2$; otherwise, let $(uu_3, uv) \to (2, 5)$.

Next assume that $\{3, 5\} \cap C(u_1) \neq \emptyset$ and $6 \notin C(u_1)$. If $3 \notin C(u_1)$, then let $(uu_1, uu_2, uu_3, uv) \to (3, 5, 6, 2)$. So assume that $3 \in C(u_1)$, i.e., $C(u_1) = \{2, 1, 4, 3\}$. Let $uu_1 \to 6$. If G contains no $(4, 2)_{(u_2,v_2)}$-path, further let $uv \to 2$; otherwise, $(uu_3, uv) \to (2, 5)$.

Case 1.2.2 $2 \in C(u_3)$ and G contains a $(2, i)_{(u_1,u_3)}$-path for any $i \in \{1, 3\} \backslash C(u_3)$.

(1.2.2.1) Assume that G contains no $(2, 3)_{(u_1,u_2)}$-path.

If G contains no $(1, 3)_{(u_2,v_1)}$-path, then let $uu_2 \to 3$. Otherwise, G contains a $(1, 3)_{(u_2,v_1)}$-path and $1 \in C(v_1)$. If $C(v_1) = \{3, 1, 2, 5\}$, then by Case 1.1, we can break that $(1, 3)_{(u,v)}$-cycle by setting $uu_2 \to 3$ and hence break the bichromatic cycle B such that no new bichromatic cycles are produced. Thus, $\{2, 5\} \backslash C(v_1) \neq \emptyset$.

Assume that there exists $a \in \{2, 5\} \backslash (C(v_1) \cup C(v_2))$. If G contains no $(6, a)_{(v_1,v_3)}$-path, then let $(vv_1, uv) \to (a, 3)$; otherwise, let $(vv_2, uu_2, uv) \to (a, 3, 4)$. Thus, assume that $\{2, 5\} \subseteq C(v_1) \cup C(v_2)$ and $2 \in C(v_1) \backslash C(v_2)$, $5 \in C(v_2) \backslash C(v_1)$ (if $5 \in C(v_1) \backslash C(v_2)$ and $2 \in C(v_2) \backslash C(v_1)$, we can let $uu_2 \to 3$ and have a similar discussion).

If G contains neither a $(2, 3)_{(v_1,v_2)}$-path nor a $(2, 6)_{(v_2,v_3)}$-path, then let $(vv_2, uu_2, uv) \to (2, 3, 4)$. Otherwise, G contains either a $(2, 3)_{(v_1,v_2)}$-path or a $(2, 6)_{(v_2,v_3)}$-path.

- Assume that G contains a $(2, 3)_{(v_1,v_2)}$-path, implying $C(v_2) = \{4, 1, 5, 3\}$.

If $\{1, 2\} \backslash C(v_3) \neq \emptyset$, then let $uv \to 6$ and $vv_3 \to \alpha \in \{1, 2\} \backslash C(v_3)$; otherwise, $\{1, 2\} \subseteq C(v_3)$. If $4 \notin C(v_1)$ and G contains no $(4, 6)_{(v_1,v_3)}$-path, then let $(vv_1, vv_2, uv) \to (4, 2, 3)$; otherwise, $4 \in C(v_1)$, or G contains a $(4, 6)_{(v_1,v_3)}$-path and

$C(v_1) = \{3, 1, 2, 6\}$, $C(v_3) = \{6, 1, 2, 4\}$.

If $4 \notin C(v_1)$, then let $(vv_1, uv) \to (5, 3)$. Otherwise, $C(v_1) = \{3, 1, 2, 4\}$. If G contains no $(4, 5)_{(v_1,v_2)}$-path, then let $(vv_1, uv) \to (5, 3)$; otherwise, G contains a $(4, 5)_{(v_1,v_2)}$-path. If $5 \notin C(v_3)$, then let $(vv_3, uv) \to (5, 6)$. Otherwise, $C(v_3) = \{6, 1, 2, 5\}$ and let $\{uu_2, vv_3\} \to 3$, $(vv_1, vv_2, uv) \to (5, 6, 4)$.

- Assume that G contains a $(2, 6)_{(v_2,v_3)}$-path, implying $C(v_2) = \{4, 1, 5, 6\}$.

First, assume that $C(v_1) = \{3, 1, 2, 4\}$. If G contains no $(4, 5)_{(v_1,v_2)}$-path, let $(vv_1, uv) \to (5, 3)$; otherwise, G contains a $(4, 5)_{(v_1,v_2)}$-path. If $5 \notin C(v_3)$, let $(vv_3, uu_2, uv) \to (5, 3, 6)$. If $3 \notin C(v_3)$, let $\{uu_2, vv_3\} \to 3$ and $(vv_2, vv_1, uv) \to (2, 6, 4)$. Otherwise, $C(v_3) = \{6, 2, 5, 3\}$ and let $(vv_1, vv_2, vv_3, uv) \to (6, 2, 4, 3)$.

Next, assume that $C(v_1) = \{3, 1, 2, 6\}$. If $3 \notin C(v_3)$, then let $\{vv_3, uu_2\} \to 3$ and $(vv_1, vv_2, uv) \to (4, 2, 6)$. If $4 \notin C(v_3)$, then let $(vv_2, vv_3, uv) \to (2, 4, 6)$. Otherwise, $C(v_3) = \{6, 2, 3, 4\}$ and let $(vv_1, vv_3, uv) \to (5, 1, 3)$.

(1.2.2.2) Assume that G contains a $(2, 3)_{(u_1,u_2)}$-path.

Then $3 \in C(u_3)$ and $1 \in C(u_1) \cup C(u_3)$ by $(*_{\mathbf{1.2}})$. If $4 \notin C(u_1) \cup C(u_3)$, let $(uu_1, uu_2, uv) \to (4, 3, 2)$. Otherwise, $4 \in C(u_1) \cup C(u_2)$. If $4 \notin C(u_3)$ and $6 \notin C(u_1) \cup C(u_3)$, let $(uu_3, uv) \to (6, 5)$. Otherwise, $4 \in C(u_3)$ or $6 \in C(u_1) \cup C(u_3)$.

- Assume that $C(u_3) = \{5, 2, 3, 1\}$ and $C(u_1) = \{2, 3, 6, 4\}$.

If G contains no $(4, 2)_{(u_2,v_2)}$-path, let $(uu_1, uv) \to (1, 2)$; otherwise, let $(uu_1, uu_2, uu_3, uv) \to (5, 3, 4, 2)$.

- Assume that $C(u_3) = \{5, 2, 3, 6\}$ and $C(u_1) = \{2, 3, 1, 4\}$.

If G contains no $(4, 2)_{(u_2,v_2)}$-path, let $(uu_1, uu_3, uv) \to (5, 1, 2)$; otherwise, let $(uu_1, uu_2, uu_3, uv) \to (5, 3, 4, 2)$.

- Assume that $C(u_3) = \{5, 2, 3, 4\}$ and $\{2, 3, 1\} \subseteq C(u_1)$.

If $4, 5 \notin C(u_1)$, let $(uu_1, uu_2, uv) \to (4, 3, 2)$. If $5, 6 \notin C(u_1)$, let $(uu_3, uu_2) \to (6, 5)$. Otherwise, $\{4, 5\} \cap C(u_1) \neq \emptyset$ and $\{5, 6\} \cap C(u_1) \neq \emptyset$. Thus, $C(u_1) = \{2, 3, 1, 5\}$. If G contains no $(5, 6)_{(u_1,v_3)}$-path, let $(uu_1, uu_3, uv) \to (6, 1, 5)$; otherwise, let $(uu_1, uu_2, uu_3, uv) \to (6, 5, 1, 2)$.

Case 1.3 $C(u_2) = \{1, 4, 2, 5\}$.

If $C(v_2) \neq \{1, 4, 3, 6\}$, then we can reduce to Cases 1.1 or 1.2. Thus, $C(v_2) = \{1, 4, 3, 6\}$. If $\{1, 3, 6\} \backslash C(u_1) \neq \emptyset$ and G contains no $(5, i)_{(u_1,u_3)}$-path for some $i \in \{1, 3, 6\} \backslash C(u_1)$, then let $(uu_1, uv) \to (i, 2)$. Otherwise, assume that:

$(*_{\mathbf{1.3}})$

(a) $\{1, 3, 6\} \subseteq C(u_1)$, or G contains a $(5, i)_{(u_1,u_3)}$-path for any $i \in \{1, 3, 6\} \backslash C(u_1)$;

(b) $\{1, 3, 6\} \subseteq C(u_3)$, or G contains a $(2, i)_{(u_1,u_3)}$-path for any $i \in \{1, 3, 6\} \backslash C(u_3)$;

(c) $\{1,2,5\} \subseteq C(v_3)$, or G contains a $(3,i)_{(v_1,v_3)}$-path for any $i \in \{1,2,5\} \backslash C(v_3)$;

(d) $\{1,2,5\} \subseteq C(v_1)$ or G contains a $(6,i)_{(v_1,v_3)}$-path for any $i \in \{1,2,5\} \backslash C(v_1)$;

(e) $\{1,3,6\} \subseteq C(u_1) \cup C(u_3)$ and $\{1,2,5\} \subseteq C(v_1) \cup C(v_3)$;

(f) $5 \in C(u_1)$ if $\{1,3,6\} \backslash C(u_1) \neq \emptyset$, $2 \in C(u_3)$ if $\{1,3,6\} \backslash C(u_3) \neq \emptyset$, $6 \in C(v_1)$ if $\{1,2,5\} \backslash C(v_1) \neq \emptyset$, and $3 \in C(v_3)$ if $\{1,2,5\} \backslash C(v_3) \neq \emptyset$.

Case 1.3.1 $5 \notin C(u_1)$.

We see that $C(u_1) = \{2,1,3,6\}$ by $(*_{\mathbf{1.3}})$. Note that G contains a $(2,i)_{(u_1,u_3)}$-path for any $i \in \{3,6\} \backslash C(u_3)$. If $\{1,3,6\} \backslash C(u_3) \neq \emptyset$, then let $(uu_1, uv) \to (5,2)$ and $uu_3 \to \beta \in \{1,3,6\} \backslash C(u_3)$. So assume that $C(u_3) = \{5,1,3,6\}$.

First assume that G contains no $(3,2)_{(u_2,v_1)}$-path. If G contains no $(3,5)_{(u_2,u_3)}$-path, let $(uu_1, uu_2, uv) \to (4,3,2)$; otherwise, let $(uu_1, uu_2, uu_3, uv) \to (5,3,4,2)$.

Next assume that G contains a $(3,2)_{(u_2,v_1)}$-path and a $(6,2)_{(u_2,v_3)}$-path. If G contains no $(1,2)_{(u_1,v_2)}$-path, let $vv_2 \to 2$; otherwise, let $\{vv_2, uu_3\} \to 2$ and $uu_1 \to 5$.

Case 1.3.2 $5 \in C(u_1)$.

Furthermore, we may assume that $2 \in C(u_3)$, $3 \in C(v_3)$, and $6 \in C(v_1)$. Since $\{1,3,6\} \subseteq C(u_1) \cup C(u_3)$ by $(*_{\mathbf{1.3}})$, we suppose that $1 \in C(u_1)$ and it suffices to consider the following two subcases by symmetry:

- $C(u_1) = \{2,5,1,3\}$, $6 \in C(u_3)$ and $4 \notin C(u_3)$.

By $(*_{\mathbf{1.3}})$, G contains a $(5,6)_{(u_1,u_3)}$-path. If $\{1,2\} \backslash C(v_3) \neq \emptyset$, let $(uu_1, uu_2) \to (4,6)$ and $uv \to \gamma \in \{1,2\} \backslash C(v_3)$. Otherwise, $C(v_3) = \{3,6,1,2\}$ and G contains a $(3,5)_{(v_1,v_3)}$-path by $(*_{\mathbf{1.3}})$, hence let $(uu_2, vv_2, uv) \to (6,5,4)$.

- $C(u_1) = \{2,5,1,3\}$ and $C(u_3) = \{5,2,6,4\}$; or $C(u_1) = \{2,5,1,4\}$ and $C(u_3) = \{5,2,3,6\}$.

By $(*_{\mathbf{1.3}})$, G contains a $(5,6)_{(u_1,u_3)}$-path, and moreover G contains a $(2,3)_{(u_1,u_3)}$-path when $C(u_1) = \{2,5,1,3\}$, a $(5,3)_{(u_1,u_3)}$-path when $C(u_1) = \{2,5,1,4\}$. If $1 \notin C(v_i)$ for some $i \in \{1,3\}$, then let $uu_2 \to c(vv_i)$. Otherwise, $1 \in C(v_1) \cap C(v_3)$. Since $\{2,5\} \subseteq C(v_1) \cup C(v_3)$ and noting that $1 \notin C(u_3)$, we have to consider the following two possibilities: If $C(v_1) = \{3,6,1,2\}$ and $C(v_3) = \{3,6,1,5\}$, let $(vv_2, vv_3) \to (5,4)$. If $C(v_1) = \{3,6,1,5\}$ and $C(v_3) = \{3,6,1,2\}$, let $(vv_1, vv_2) \to (4,5)$.

Case 2 $|C(u) \cap C(v)| = 3$, say $(uu_1, uu_3)_c = (2,5)$ and $(vv_1, vv_3)_c = (3,5)$.

If $C(u_2) = \{1,4,3,6\}$ or $C(v_2) = \{1,4,2,6\}$, then the proof can be reduced to Case 1. If G contains neither a $(4,6)_{(u_2,v)}$-path nor a $(5,6)_{(u_2,v)}$-path, then let $uv \to 6$. Otherwise, assume that:

$(*_{\mathbf{2.1}})$ G contains either a $(4,6)_{(u,v)}$-path or a $(5,6)_{(u,v)}$-path.

Suppose that we can recolor some edges in G such that $|C(u) \cap C(v)| = 2$, and no new bichromatic cycles are produced in $G - uv$. Then, by the discussion of Case 1.1, if G contains an $(i, j)_{(u,v)}$-cycle for some $i, j \in C(u) \cap C(v)$, then we can break this $(i, j)_{(u,v)}$-cycle and hence break the cycle B such that no new bichromatic cycles are produced. That is, we have the following statement:

($\circledast_{\mathbf{2}}$) If we can recolor some edges in G such that $|C(u) \cap C(v)| = 2$, and no new bichromatic cycles are produced in $G - uv$, then we are done.

If $6 \notin C(u_3)$ and G contains no $(2, 6)_{(u_1,u_3)}$-path, let $uu_3 \to 6$; if $1 \notin C(u_3)$ and G contains no $(1, 2)_{(u_1,u_3)}$-path, let $(uu_3, uv) \to (1, 6)$. Clearly, $|C(u) \cap C(v)| = 2$ and no new bichromatic cycles are produced in $G - uv$. By ($\circledast_{\mathbf{2}}$), we complete the proof. Similarly, if $6 \notin C(u_2)$ and G contains no $(2, 6)_{(u_1,u_2)}$-path, we can let $uu_2 \to 6$. Hence, assume that:

($*_{\mathbf{2.2}}$)

(a) $6 \in C(u_2)$, or G contains a $(2, 6)_{(u_1,u_2)}$-path;

(b) $6 \in C(v_2)$, or G contains a $(3, 6)_{(v_1,v_2)}$-path;

(c) $\{1, 6\} \subseteq C(u_3)$, or G contains a $(2, i)_{(u_1,u_3)}$-path for each $i \in \{1, 6\} \backslash C(u_3)$;

(d) $\{1, 6\} \subseteq C(v_3)$, or G contains a $(3, i)_{(v_1,v_3)}$-path for each $i \in \{1, 6\} \backslash C(v_3)$;

(e) $\{1, 6\} \subseteq C(u_1) \cup C(u_3)$ and $\{1, 6\} \subseteq C(v_1) \cup C(v_3)$;

(f) $2 \in C(u_3)$ if $\{1, 6\} \backslash C(u_3) \neq \emptyset$ and $3 \in C(u_3)$ if $\{1, 6\} \backslash C(v_3) \neq \emptyset$.

It follows that $\{2, 6\} \cap C(u_2) \neq \emptyset$ and $\{3, 6\} \cap C(v_2) \neq \emptyset$. Thus, $C(u_2) \backslash \{1, 4\} \in \{\{2, 3\}, \{2, 5\}, \{2, 6\}, \{5, 6\}\}$ and $C(v_2) \backslash \{1, 4\} \in \{\{2, 3\}, \{3, 5\}, \{3, 6\}, \{5, 6\}\}$.

Case 2.1 $C(u_2) \backslash \{1, 4\} = \{2, 3\}$.

By ($*_{\mathbf{2.1}}$) and ($*_{\mathbf{2.2}}$), G contains a $(5, 6)_{(u,v)}$-path with $6 \in C(u_3) \cap C(v_3)$, or a $(2, 6)_{(u_1,u_2)}$-path with $6 \in C(u_1)$ and $1 \in C(u_3)$, or a $(1, 2)_{(u_1,u_3)}$-path with $2 \in C(u_3)$ and $1 \in C(u_1)$. If $4 \notin C(u_1)$ and G contains no $(4, 5)_{(u_1,u_3)}$-path, let $(uu_1, uu_2) \to (4, 6)$. Otherwise, $4 \in C(u_1)$, or G contains a $(4, 5)_{(u_1,u_3)}$-path and $5 \in C(u_1)$, $4 \in C(u_3)$. We need to consider the following subcases.

Case 2.1.1 $4 \in C(u_1)$ and $1 \in C(u_3)$.

- Assume that $C(u_1) = \{2, 6, 4, 3\}$.

If $2 \notin C(u_3)$, let $(uu_1, uu_2, uu_3, uv) \to (1, 5, 2, 6)$; otherwise, $C(u_3) = \{5, 6, 1, 2\}$, let $(uu_1, uu_2, uu_3) \to (5, 6, 4)$.

- Assume that $3 \notin C(u_1)$.

If $2 \notin C(u_3)$ and G contains no $(2,4)_{(u_2,u_3)}$-path, let $(uu_3, uv) \to (2,6)$ and $uu_1 \to \alpha \in \{1,5\} \backslash C(u_1)$. By $(*_{\mathbf{2.1}})$ and Lemma 2, no new bichromatic cycles are produced. Otherwise, $2 \in C(u_3)$ or G contains a $(2,4)_{(u_2,u_3)}$-path and $4 \in C(u_3)$.

(i) Assume that $C(u_3) = \{5,1,6,4\}$ and G contains a $(2,4)_{(u_2,u_3)}$-path.

If G contains no $(4,3)_{(u_1,u_2)}$-path, let $(uu_1, uv) \to (3,2)$; otherwise, let $(uu_3, uv) \to (3,2)$, $uu_1 \to \beta \in \{1,5\} \backslash C(u_1)$.

(ii) Assume that $C(u_3) = \{5,6,1,2\}$.

If G contains no $(3,6)_{(u_3,v_1)}$-path, let $(uu_3, uv) \to (3,6)$. So assume that G contains a $(3,6)_{(u_3,v_1)}$-path. It follows that $6 \in C(v_2)$ by $(*_{\mathbf{2.2}})$ and the fact that $2 \notin C(v_2)$. If $C(u_1) = \{2,6,4,1\}$, let $(uu_1, uu_2, uu_3, uv) \to (5,6,4,2)$. Otherwise, we derive that $C(u_1) = \{2,6,4,5\}$. If G contains no $(2,3)_{(u_3,v_1)}$-path, then let $(uu_1, uu_3, uv) \to (1,3,2)$. Thus, assume that G contains a $(2,3)_{(u_3,v_1)}$-path and $2 \in C(v_1)$. If G contains no $(2,5)_{(v_2,v_3)}$-path, then let $vv_2 \to 2$; otherwise, G contains a $(2,5)_{(v_2,v_3)}$-path, which implies that $C(v_2) = \{4,1,6,5\}$ and $2 \in C(v_3)$. If $1, 5 \notin C(v_1)$, let $(vv_1, uv) \to (1,3)$. Otherwise, $\{1,5\} \cap C(v_1) \neq \emptyset$ and then $4 \notin C(v_1)$ since $\{2,6\} \subseteq C(v_1)$. If $4 \notin C(v_3)$, then let $(vv_3, vv_2) \to (4,2)$; otherwise, let $(vv_1, vv_2, vv_3) \to (4,2,3)$.

Case 2.1.2 $4 \notin C(u_1)$ or $1 \notin C(u_3)$.

If $4 \notin C(u_1)$ and $1 \in C(u_3)$, then $C(u_3) = \{5,6,1,4\}$, let $(uu_1, uu_2, uu_3) \to (4,6,2)$. If $4 \in C(u_1)$ and $1 \notin C(u_3)$, then $C(u_1) = \{2,6,1,4\}$, let $(uu_1, uu_3, uv) \to (5,1,6)$. Otherwise, assume that $4 \notin C(u_1)$ and $1 \notin C(u_3)$. Then $C(u_1) = \{2,6,1,5\}$ and $C(u_3) = \{5,6,2,4\}$. If G contains no $(3,6)_{(u_1,v_1)}$-path, let $(uu_1, uu_3, uv) \to (3,1,6)$. Otherwise, it follows that $6 \in C(v_2)$ by $(*_{\mathbf{2.2}})$ and $2 \notin C(v_2)$. Let $(uu_1, uu_3, uv) \to (3,1,2)$.

Case 2.2 $C(u_2) \backslash \{1,4\} = \{2,5\}$.

If $C(v_2) \backslash \{1,4\} = \{2,3\}$, then the proof is reduced to Case 2.1. Thus, $C(v_2) \backslash \{1,4\} \neq \{2,3\}$ and $2 \notin C(v_2)$. By $(*_{\mathbf{2.1}})$ and $(*_{\mathbf{2.2}})$, G contains a $(5,6)_{(u_3,v_3)}$-path with $6 \in C(u_3) \cap C(v_3)$, or G contains a $(2,6)_{(u_1,u_2)}$-path with $6 \in C(u_1)$ and $1 \in C(u_3)$, or a $(1,2)_{(u_1,u_3)}$-path with $1 \in C(u_1)$ and $2 \in C(u_3)$. If $4 \notin C(u_1)$ and G contains no $(4,5)_{(u_1,u_3)}$-path, then let $(uu_1, uu_2) \to (4,6)$. Otherwise, $4 \in C(u_1)$, or G contains no $(4,5)_{(u_1,u_3)}$-path and $5 \in C(u_1)$, $4 \in C(u_3)$. We need to consider the following two subcases.

Case 2.2.1 $4 \in C(u_1)$ and $1 \in C(u_3)$.

- Assume that $C(u_1) = \{2,6,4,1\}$.

If $2 \notin C(u_3)$, let $(uu_1, uv) \to (3,2)$. Otherwise, $C(u_3) = \{5,6,1,2\}$, let $(uu_1, uu_2, uu_3, uv) \to (3,6,4,2)$.

- Assume that $1 \notin C(u_1)$.

If $2,4 \notin C(u_3)$, let $(uu_1, uu_3, uv) \to (1,2,6)$. Otherwise, $\{2,4\} \cap C(u_3) \neq \emptyset$.

(i) Assume that $C(u_3) = \{5,1,6,4\}$.

If $5 \notin C(u_1)$, let $(uu_1, uv) \to (1,2)$. Otherwise, $C(u_1) = \{2,6,4,5\}$, we let $(uu_1, uv) \to (3,2)$.

(ii) Assume that $C(u_3) = \{5,6,1,2\}$.

If $3 \notin C(u_1)$, let $(uu_1, uu_2) \to (3,6)$. Otherwise, $C(u_1) = \{2,6,4,3\}$.

If G contains no $(2,5)_{(u_3,v_3)}$-path, let $(uu_1, uv) \to (1,2)$. So assume that G contains a $(2,5)_{(u_3,v_3)}$-path and $2 \in C(v_3)$. If G contains no $(2,3)_{(v_1,v_2)}$-path, let $vv_2 \to 2$. Thus, assume that G contains a $(2,3)_{(v_1,v_2)}$-path and $2 \in C(v_1)$, $3 \in C(v_2)$. Note that $1 \in C(v_3)$ or G contains a $(1,3)_{(v_1,v_3)}$-path, and $\{1,3\} \cap C(v_3) \neq \emptyset$ by $(*_{\mathbf{2.2}})$ and then $4 \notin C(v_3)$. If $4 \notin C(v_1)$, then let $(vv_1, vv_2) \to (4,2)$. Otherwise, $4 \in C(v_1)$.

Assume that $1 \notin C(v_3)$ and then $C(v_3) = \{5,6,2,3\}$, $C(v_1) = \{3,2,4,1\}$ by $(*_{\mathbf{2.2}})$. If $6 \notin C(v_2)$, let $vv_2 \to 6$; otherwise, $C(v_2) = \{4,1,3,6\}$, let $(vv_1, vv_3, uv) \to (5,1,6)$. Next, assume that $C(v_3) = \{5,6,2,1\}$. If $1 \notin C(v_1)$, let $(vv_1, vv_3, uv) \to (1,3,6)$; otherwise, $C(v_1) = \{3,2,4,1\}$. If $6 \notin C(v_2)$, then let $vv_2 \to 6$; otherwise, $C(v_2) = \{4,1,3,6\}$, let $(vv_1, vv_3, uv) \to (5,3,6)$.

Case 2.2.2 $4 \notin C(u_1)$ or $1 \notin C(u_3)$.

If $4 \notin C(u_1)$ and $1 \in C(u_3)$, then $C(u_3) = \{5,6,1,4\}$, let $(uu_1, uu_2, uu_3) \to (4,6,2)$. If $4 \notin C(u_1)$ and $1 \notin C(u_3)$, then $C(u_1) = \{2,6,1,5\}$, $C(u_3) = \{5,6,2,4\}$, and let $(uu_1, uu_3, uv) \to (3,1,2)$. Otherwise, assume that $4 \in C(u_1)$ and $1 \notin C(u_3)$. Then $C(u_1) = \{2,6,1,4\}$. If G contains no $(1,3)_{(u_1,u_3)}$-path, then let $(uu_1, uu_3, uv) \to (3,1,2)$; otherwise, let $(uu_1, uu_2, uu_3, uv) \to (5,3,1,6)$.

Case 2.3 $C(u_2) \backslash \{1,4\} = \{2,6\}$.

If $C(v_2) \backslash \{1,4\} \in \{\{2,3\},\{3,5\}\}$, then we reduce the proof to Cases 2.1 or 2.2. Thus, $C(v_2) \backslash \{1,4\} \in \{\{3,6\},\{5,6\}\}$.

Case 2.3.1 G contains no $(2,3)_{(u_1,u_2)}$-path.

First let $uu_2 \to 3$. If G contains no $(1,3)_{(u_2,v_2)}$-path, then we are done. Otherwise, G contains a $(1,3)_{(u_2,v_2)}$-path and $|C(u) \cap C(v)| = 2$.

If $C(v_1) \backslash \{1,3\} \notin \{\{4,6\},\{5,6\}\}$, then by Case 1, Case 2.1 or Case 2.2, we can break this $(1,3)_{(u,v)}$-cycle and hence break the cycle B and no new bichromatic cycles are produced. Thus, $C(v_1) \backslash \{1,3\} \in \{\{4,6\},\{5,6\}\}$.

Note that G contains a $(3,i)_{(v_1,v_3)}$-path for any $i \in \{1,6\} \backslash C(v_3)$ by $(*_{\mathbf{2.2}})$. This implies that $1 \in C(v_3)$. If $6 \notin C(v_3)$, let $uv \to 6$. Hence, $\{1,6\} \subseteq C(v_3)$.

- Assume that $C(v_2) = \{4, 1, 6, 3\}$. (If $C(v_1) = \{3, 1, 6, 4\}$, we have a similar discussion.)

If $4 \notin C(v_3)$, let $(vv_2, uv) \to (2, 4)$. Otherwise, $C(v_3) = \{5, 1, 6, 4\}$, we let $(vv_1, uu_2, uv) \to (2, 4, 3)$.

- Assume that $C(v_2) = \{4, 1, 6, 5\}$ and $C(v_1) = \{3, 1, 6, 5\}$.

If $3, 4 \notin C(v_3)$, let $(vv_1, vv_2) = (4, 3)$. Otherwise, $\{3, 4\} \cap C(v_3) \neq \emptyset$. If $C(v_3) = \{5, 1, 6, 3\}$, let $(vv_2, uv) = (2, 4)$; otherwise, $C(v_3) = \{5, 1, 6, 4\}$, let $(uu_2, vv_1, uv) = (4, 2, 3)$.

Case 2.3.2 G contains a $(2, 3)_{(u_1,u_2)}$-path and $3 \in C(u_1)$.

(2.3.2.1) $C(v_2) = \{4, 1, 5, 6\}$.

- Assume that G contains no $(2, 5)_{(v_2,v_3)}$-path. Let $vv_2 \to 2$. If G contains no $(1, 2)_{(u,v)}$-path, then we are done; otherwise, G contains a $(1, 2)_{(u,v)}$-path, we can break this $(1, 2)_{(u,v)}$-cycle by Case 1 or Case 2.1 as $3 \in C(u_1)$.
- Assume that G contains a $(2, 5)_{(v_2,v_3)}$-path. If $2 \notin C(v_1)$ and G contains no $(5, 3)_{(u_3,v_3)}$-path, then let $(vv_1, uv) \to (2, 3)$. So assume that $2 \in C(v_1)$ or G contains a $(5, 3)_{(u_3,v_3)}$-path. If $2 \notin C(v_1)$ and G contains a $(5, 3)_{(u_3,v_3)}$-path, let $(vv_1, vv_2) \to (2, 3)$. If $4 \notin C(v_3)$ and G contains no $(3, 4)_{(v_1,v_3)}$-path, let $(vv_3, vv_2) \to (4, 2)$. If G contains no $(1, 2)_{(u,v)}$-path, we are done; otherwise, G contains a $(1, 2)_{(u,v)}$-path, we can break this $(1, 2)_{(u,v)}$-cycle by Cases 1 or 2.1 as $3 \in C(u_1)$. Otherwise, $2 \in C(v_1)$, and $4 \in C(v_3)$ or G contains a $(3, 4)_{(v_1,v_3)}$-path. Together with $(*_{\mathbf{2.2}})$, we have $3 \in C(v_3)$, $\{1, 6\} \backslash C(v_3) \neq \emptyset$, and then $5 \notin C(v_1)$ since $\{1, 6, 4\} \subseteq C(v_1) \cup C(v_3)$. Let $(vv_1, vv_2) \to (5, 3)$, $vv_3 \to \beta \in \{1, 6\} \backslash C(v_1)$, and $uv \to \{1, 6\} \backslash \beta$. By $(\divideontimes_{\mathbf{2}})$, we complete the proof.

(2.3.2.2) $C(v_2) = \{4, 1, 3, 6\}$.

Similarly, we can assume that G contains a $(2, 3)_{(v_1,v_2)}$-path and $2 \in C(v_1)$.

First, assume that $2 \notin C(u_3)$, and hence $\{1, 6\} \subseteq C(u_3)$ by $(*_{\mathbf{2.2}})$. If $4 \notin C(u_1) \cup C(u_3)$, let $(uu_1, uu_2, uv) \to (4, 3, 2)$; otherwise, $4 \in C(u_1) \cup C(u_3)$. If $1 \notin C(u_1)$ and G contains no $(2, 4)_{(u_2,u_3)}$-path, then let $(uu_1, uu_3, uv) \to (1, 2, 6)$ and we are done by $(\divideontimes_{\mathbf{2}})$; otherwise, $1 \in C(u_1)$ or G contains a $(2, 4)_{(u_2,u_3)}$-path. Assume that $C(u_3) = \{5, 1, 6, 3\}$ and then $\{1, 4\} \subseteq C(u_1)$ since G contains no $(2, 4)_{(u_2,u_3)}$-path and $4 \in C(u_1) \cup C(u_3)$. Then, let $(uu_1, uu_2, uu_3, uv) \to (5, 3, 4, 2)$. Otherwise, assume that $C(u_3) = \{5, 1, 6, 4\}$. If $5 \notin C(u_1)$, let $(uu_1, uu_3, uv) \to (5, 3, 2)$; otherwise, $5 \in C(u_1)$. Assume that $1 \notin C(u_1)$. If G contains no $(1, 5)_{(u_1,u_3)}$-path, let $(uu_1, uv) \to (1, 2)$; otherwise, G contains a $(1, 5)_{(u_1,u_3)}$-path, let $(uu_1, uu_2, uu_3, uv) \to (1, 5, 2, 6)$ and we are done by $(\divideontimes_{\mathbf{2}})$. Otherwise, $C(u_1) = \{2, 3, 5, 1\}$. If G contains no $(5, 6)_{(u_1,u_3)}$-path, then let $(uu_1, uv) \to (6, 2)$; otherwise, G contains a $(5, 6)_{(u_1,u_3)}$-path, let $(uu_1, uu_2, uu_3) \to (6, 5, 2)$ and we are done by $(\divideontimes_{\mathbf{2}})$.

Next, assume that $2 \in C(u_3)$ and $3 \in C(v_3)$. Assume that $4 \notin C(u_3)$. If $5 \notin C(u_1)$, let $(uu_1, uu_2, uu_3, uv) \to (5, 3, 4, 2)$. So assume that $5 \in C(u_1)$.

Assume that $4 \notin C(u_1)$. Note that G contains a $(4, 6)_{(u_2,v_2)}$-path and a $(3, i)_{(v_1,v_3)}$-path for any $i \in \{1, 6\}\backslash C(v_3)$ by $(*_{\mathbf{2.1}})$ and $(*_{\mathbf{2.2}})$. If $\{1, 6\}\backslash C(v_3) \neq \emptyset$, then let $(uu_1, uu_2) \to (4, 3)$ and $uv \to \alpha \in \{1, 6\}\backslash C(v_3)$. Otherwise, $C(v_3) = \{5, 3, 1, 6\}$, let $(uu_1,\ uu_2,\ uv) \to (4, 3, 2)$.

It remains to consider the case $C(u_1) = \{2, 3, 5, 4\}$ and $C(u_3) = \{5, 2, 6, 1\}$ since $\{1, 6\} \subseteq C(u_1) \cup C(u_3)$ by $(*_{\mathbf{2.2}})$. If $2 \notin C(v_3)$, let $(vv_3, uu_3, uv) \to (2, 3, 5)$. So assume that $2 \in C(v_3)$. If $1, 6 \notin C(v_3)$, then $C(v_1) = \{3, 2, 1, 6\}$, let $(vv_1, vv_2) \to (4, 2)$. Otherwise, $\{1, 6\} \cap C(v_3) \neq \emptyset$ and $4 \notin C(v_3)$. If $4 \notin C(v_1)$, let $(vv_1, vv_2) \to (4, 2)$; otherwise, $4 \in C(v_1)$ and $5 \notin C(v_1)$, let $(vv_1, vv_2, vv_3, uv) \to (5, 2, 4, 3)$.

Hence, $4 \in C(u_3) \cap C(v_3)$. If $3 \notin C(u_3)$ and $2 \notin C(v_3)$, let $(uu_3,\ vv_3,\ uv) \to (3, 2, 5)$. Otherwise, $3 \in C(u_3)$ or $2 \in C(v_3)$ and assume that $C(u_3) = \{5, 2, 4, 3\}$ and $C(u_1) = \{3, 2, 1, 6\}$ by $(*_{\mathbf{2.2}})$. Let $(uu_1, uu_3, uv) \to (5, 1, 2)$.

Case 2.4 $C(u_2)\backslash\{1, 4\} = \{5, 6\}$.

If $C(v_2)\backslash\{1, 4\} \neq \{5, 6\}$, then we reduce the proof to Cases 2.1-2.3. Thus, $C(v_2)\backslash\{1, 4\} = \{5, 6\}$.

Case 2.4.1 G contains no $(5, 3)_{(u_2,u_3)}$-path.

First, let $uu_2 \to 3$. If G contains no $(1, 3)_{(u_2,v_2)}$-path, we are done. Otherwise, G contains a $(1, 3)_{(u_2,v_2)}$-path and $1 \in C(v_3)$ by $(*_{\mathbf{2.2}})$. If $C(v_1)\backslash\{1, 3\} \neq \{5, 6\}$, then by Cases 1, 2.1-2.3, we can break this $(1, 3)_{(u,v)}$-cycle and hence break the cycle B so that no new bichromatic cycles are produced. Thus, assume that $C(v_1)\backslash\{1, 3\} = \{5, 6\}$. Note that G contains a $(4, 6)_{(u_2,v_2)}$-path and a $(3, 6)_{(v_1,v_3)}$-path by $(*_{\mathbf{2.1}})$ and $(*_{\mathbf{2.2}})$. If $6 \notin C(v_3)$, let $uv \to 6$. Otherwise, $6 \in C(v_3)$. If $3, 4 \notin C(v_3)$, let $(vv_1, vv_2) = (4, 3)$; otherwise, $\{3, 4\} \cap C(v_3) \neq \emptyset$. If $C(v_3) = \{5, 1, 6, 3\}$, let $(vv_2, uv) = (2, 4)$; otherwise, $C(v_3) = \{5, 1, 6, 4\}$, assuming that $(uu_2, vv_1, uv) = (4, 2, 3)$, $C(v_3) = \{5, 1, 6, 3\}$, let $(vv_2, uv) \to (2, 4)$.

Case 2.4.2 G contains a $(3, 5)_{(u_2,u_3)}$-path and $(2, 5)_{(v_2,v_3)}$-path.

If $2 \notin C(v_1)$, let $(vv_1, uv) \to (2, 3)$; otherwise, $2 \in C(v_1)$. Note that $\{1, 6\} \subseteq C(v_3)$ or G contains a $(3, i)_{(v_1,v_3)}$-path for any $i \in \{1, 6\}\backslash C(v_3)$ by $(*_{\mathbf{2.2}})$.

Assume that $3 \notin C(v_3)$ and then $C(v_3) = \{5, 2, 1, 6\}$ by $(*_{\mathbf{2.2}})$. If $4 \notin C(v_1)$, let $(vv_1, vv_2) \to (4, 3)$. If $1 \notin C(v_1)$, let $(vv_1, vv_3, uv) \to (1, 3, 6)$ and we are done by $(\divideontimes_{\mathbf{2}})$. Otherwise, $C(v_1) = \{3, 2, 4, 1\}$ and let $(vv_1, vv_2) \to (6, 3)$.

Assume that $3 \in C(v_3)$ and $\{1,6\}\backslash C(v_3) \neq \emptyset$. If $5 \notin C(v_1)$, let $(vv_1, vv_2) \rightarrow (5,3)$, $vv_3 \rightarrow \beta_1 \in \{1,6\}\backslash C(v_3)$, $uv \rightarrow \{1,6\}\backslash \beta_1$. Otherwise, $5 \in C(v_1)$. Since $\{1,6\} \subseteq C(v_1) \cup C(v_3)$ by $(*_{\mathbf{2.2}})$, we have $4 \notin C(v_1) \cup C(v_3)$. Let $(vv_1, vv_2) \rightarrow (4,3)$, $vv_3 \rightarrow \beta_2 \in \{1,6\}\backslash C(v_3)$, $uv \rightarrow \{1,6\}\backslash \beta_2$.

Case 3 $|C(u) \cap C(v)| = 4$, say $(uu_1, uu_3)_c = (2,5)$ and $(vv_1, vv_3)_c = (2,5)$.

If $C(u_2)\backslash \{1,4\} \neq \{2,5\}$ or $C(v_2)\backslash\{1,4\} \neq \{2,5\}$, then $|C(u_2) \cap C(u)| \leq 3$ or $|C(v_2) \cap C(v)| \leq 3$ and we reduce the proof to Cases 1 or 2. Thus, $C(u_2) = C(v_2) = \{1,2,4,5\}$. If G contains neither a $(2,j)_{(u,v)}$-path nor a $(5,j)_{(u,v)}$-path for some $j \in \{3,6\}$, then let $uv \rightarrow j$. Otherwise, for any $j \in \{3,6\}$, G contains an $(i,j)_{(u,v)}$-path for some $i \in \{2,5\}$, and G contains a $(2,3)_{(u,v)}$-path, $3 \in C(u_1)$. If G contains no $(3,5)_{(v_2,v_3)}$-path, let $vv_2 \rightarrow 3$. Otherwise, G contains a $(3,5)_{(v_2,v_3)}$-path and a $(3,5)_{(u_2,u_3)}$-path similarly. It follows that $3 \in C(u_3)$ and $6 \in C(u_1) \cap C(u_3)$ similarly. If $1 \notin C(u_1)$ and G contains no $(1,5)_{(u_1,u_3)}$-path, then let $(uu_1, uv) \rightarrow (1,3)$. Hence, $1 \in C(u_1)$ or G contains a $(1,5)_{(u_1,u_3)}$-path. It follows that $\{1,5\} \cap C(u_1) \neq \emptyset$, $4 \notin C(u_1)$, and if $1 \notin C(u_1)$, then $1 \in C(u_3)$. If $4 \notin C(u_3)$, let $(uu_2, uu_3) \rightarrow (3,4)$. Otherwise, $C(u_3) = \{5,3,6,4\}$ and $C(u_1) = \{2,3,6,1\}$. Let $(uu_1, uu_3, uv) \rightarrow (5,1,3)$. □

Lemma 3 *([4]) If G is a graph with $\Delta = 4$ that is not 4-regular, then $a'(G) \leq 6$.*

Theorem 2 *If G is a 4-regular graph, then $a'(G) \leq 6$.*

Proof. The proof is proceeded by induction on the vertex number of G. If $|V(G)| = 5$, then G is the complete graph K_5, and it is easy to show that $a'(G) \leq 6$. Let G be a 4-regular graph with $|V(G)| \geq 6$. Obviously we may assume that G is 2-connected by Lemma 3. If G contains no 3-cycles, then $a'(G) \leq 6$ by Theorem 1. So assume that G contains at least one 3-cycle. For any graph H with $\Delta(H) = 4$ and $|V(H)| < |V(G)|$, by induction hypothesis or Lemma 3, H admits an acyclic edge-6-coloring c using the color set $C = \{1, 2, \dots, 6\}$. Let $v \in V(G)$ be a vertex with neighbors v_0, v_1, v_2, v_3 that lies in at least one 3-cycle. To extend c to the original graph G, we split the proof into the following cases.

Case 1 $v_1v_2, v_2v_3, v_3v_0, v_0v_1 \in E(G)$, as shown in Fig. 3.

For $i \in \{0,1,2,3\}$, let v_i' be the neighbor of v_i different from v, v_{i-1}, v_{i+1}, where the indices are taken modulo 4. Since $|V(G)| \geq 6$, $G \neq K_5$. We only need to consider the following subcases by symmetry:

Case 1.1 $v_1v_3 \in E(G)$ and $v_2v_0 \notin E(G)$.



Case 1.1

Case 1.2

Fig. 3: The configuration in Case 1.

Since G is 2-connected, $v_0' \neq v_2'$. Let $H = G - \{v, v_1, v_2, v_3, v_0\} + \{u, uv_2', uv_0'\}$. Without loss of generality, assume that $c(uv_2') = 1$ and $c(uv_0') = 2$. To extend c to the graph G, we let $\{v_1v_3, v_2v_2', vv_0\} \to 1$, $\{vv_3, v_0v_0'\} \to 2$, $\{v_1v_0, v_2v_3\} \to 5$, $\{v_3v_0, vv_2\} \to 6$, and $(vv_1, v_1v_2) \to (3, 4)$.

Case 1.2 $v_1v_3, v_2v_0 \notin E(G)$.

If v_0', v_1', v_2', v_3' are identical into a vertex, then $|V(G)| = 6$ and it is easy to give an acyclic edge-6-coloring of G.

If v_1', v_2', v_3' are identical to a vertex, say v_5, and $v_0' \neq v_5$, then let $H = G - \{v, v_1, v_2, v_3, v_0\} + \{u, uv_0', uv_5\}$ with $(uv_5, uv_0')_c = (1, 4)$, $2, 3 \notin C(v_5)$, where u is a new vertex added in the graph.

If v_1', v_2' are are identical to a vertex v_5 and v_0', v_3' are are identical into a vertex v_6, we let $H = G - \{v, v_1, v_2, v_3, v_0\} + \{u, uv_5, uv_6\}$. Assume that $(uv_5, uv_6)_c = (1, 4)$, $2 \in C \backslash (C(v_5) \cup \{1, 4\})$, and $3 \in C \backslash (C(v_6) \cup \{1, 4, 2\})$.

If v_1', v_2' are are identical to a vertex v_5, and $v_3', v_0' \neq v_5$, let $H = G - \{v, v_1, v_2, v_3, v_0\} + \{u, uv_5, uv_3', uv_0'\}$ and assume that $(uv_5, uv_3', uv_0')_c = (1, 3, 4)$, $2 \in C \backslash (C(v_5) \cup \{3, 4\})$.

If $v_1' = v_3' = v_5$ and $v_2' = v_0' = v_6$, let $H = G - \{v, v_1, v_2, v_3, v_0\} + \{u, uv_5, uv_6\}$ and assume that $(uv_5, uv_6)_c = (1, 2)$, $3 \in C \backslash (C(v_5) \cup \{1, 2\})$, and $4 \in C \backslash (C(v_6) \cup \{1, 2, 3\})$.

If $v_1' = v_3' = v_5$ and $v_2' \neq v_0' \neq v_5$, let $H = G - \{v, v_1, v_2, v_3, v_0\} + \{u, uv_5, uv_2', uv_0'\}$ and assume that $(uv_5, uv_2', uv_0')_c = (1, 2, 4)$, $3 \in C \backslash (C(v_5) \cup \{1, 2\})$.

If v_0', v_1', v_2', v_3' are all distinct, let $H = G - \{v, v_1, v_2, v_3, v_0\} + \{u, uv_1', uv_2', uv_3', uv_0'\}$ and assume that $(uv_1', uv_2', uv_3', uv_0')_c = (1, 2, 3, 4)$.

For all cases above, we first let $v_iv_i' \to i$ for $i = 1, 2, 3$ and $v_0v_0' \to 4$. Then let $\{vv_2, v_1v_0\} \to 5$, $\{vv_0, v_2v_3\} \to 6$, and $(v_3v_0, vv_3, v_1v_2, vv_1) \to (1, 2, 3, 4)$.

Case 2 $v_0v_1, v_1v_2, v_2v_3 \in E(G)$ and $v_3v_0 \notin E(G)$, as shown in Fig. 4.



Case 2.1

Case 2.2

Fig. 4: The configuration in Case 2.

Let v_2' be the neighbor of v_2 different from v, v_1, v_3. If $v_1v_3, v_2v_0 \in E(G)$, then $v_1v_0, v_0v_2, v_2v_3, v_3v_1 \in E(G)$ and the proof is reduced to Case 1. Without loss of generality, assume that $v_2v_0 \notin E(G)$.

Case 2.1 $v_1v_3 \in E(G)$.

Let v_3' be the neighbor of v_3 different from v, v_1, v_2.

If v_2', v_3' are identical to a vertex, say v_5, then let $H = G - \{v, v_1, v_2, v_3\} + \{u, uv_5, uv_0\}$ and assume that $(uv_5, uv_0)_c = (1, 2)$, $5 \in C\backslash(C(v_5) \cup \{1, 2\})$, and $6 \in C\backslash(C(v_0) \cup \{1, 2, 5\})$.

If $v_3' \neq v_2'$, let $H = G - \{v, v_1, v_2, v_3\} + \{u, uv_0, uv_2', uv_3'\}$ and assume that $(uv_0, uv_2', uv_3')_c = (2, 5, 1)$ and $6 \in C\backslash(C(v_0) \cup \{1, 2, 5\})$.

For two cases above, we further let $\{v_1v_2, v_3v_3'\} \to 1$, $\{v_2v_3, v_1v_0\} \to 2$, $\{v_2v_5, vv_3\} \to 5$, $\{vv_0, v_1v_3\} \to 6$, and $(vv_2, vv_1) \to (3, 4)$.

Case 2.2 $v_1v_3 \notin E(G)$.

Let v_5 be the neighbor of v_1 different from v, v_0, v_2. We consider two subcases as follows:

Case 2.2.1 $v_2' = v_5$.

By Case 1, $v_5v_3, v_5v_0 \notin E(G)$. Let $H = G - \{v, v_1, v_2\} + \{v_5v_3, v_5v_0\}$ and assume that $(v_5v_3, v_5v_0)_c = (2, 1)$. Then, first, let $\{v_1v_5, vv_0\} \to 1$, $\{v_2v_5, vv_3\} \to 2$. Next, if $\{3, 4, 5, 6\}\backslash(C(v_3) \cup C(v_0)) \neq \emptyset$ and assume that $3 \notin C(v_3) \cup C(v_0)$, $4 \notin C(v_0)$, then let $\{v_1v_0, v_2v_3\} \to 3$ and $(v_1v_2, vv_1, vv_2) \to (4, 5, 6)$. Otherwise, $C(v_0) = \{3, 4\}$ and $C(v_3) = \{5, 6\}$. Then, let $\{v_1v_0, vv_2\} \to 6$ and $(v_1v_2, v_2v_3, vv_1) \to (3, 4, 5)$. Since H has no bichromatic cycles, it is obvious that H has no $(1, 2)_{(v_0, v_3)}$-path and thus G

contains no $(1,2)_{(v_0,v)}$-cycle. (See Fig. 5. We neglect to explain why such bichromatic cycles cannot exist in the following similar cases.)



Fig. 5: Case 2.2.1.

Case 2.2.2 $v_2' \neq v_5$.

Let $H = G - \{v, v_1, v_2\} + \{u, uv_5, uv_2', uv_3, uv_0, v_3v_0\}$ and assume that $(uv_0, uv_3, v_3v_0)_c = (1,2,3)$. Then, first let $(v_1v_5, v_2v_2') \to (c(uv_5), c(uv_2'))$.

- Assume that $(uv_5, uv_2') = (3,4)$.

If $\{5,6\} \backslash C(v_3) \neq \emptyset$ and $5 \notin C(v_3)$, then let $\{v_1v_0, vv_2\} \to 1$, $\{vv_1, v_2v_3\} \to 2$, and $(vv_0, vv_3, v_1v_2) \to (3,5,6)$. Otherwise, $C(v_3) = \{5,6\}$. Then, first let $\{vv_0, v_1v_2\} \to 1$, $(vv_3, v_2v_3) \to (2,3)$. And next, if $4 \notin C(v_0)$, then let $(v_1v_0, vv_1, vv_2) \to (4,5,6)$. Otherwise, $4 \in C(v_0)$ and assume that $5 \notin C(v_0)$, then let $(vv_1, v_1v_0, vv_2) \to (4,5,6)$.

- Assume that $(uv_5, uv_2') = (4,5)$.

If $4 \notin C(v_0)$, then let $\{vv_0, v_1v_2\} \to 1$, $\{v_1v_0, vv_3\} \to 3$, and $(v_2v_3, vv_2, vv_1) \to (2,4,6)$. Since H has no bichromatic cycles and $c(v_0v_3) = 3$, it is obvious that H has no $(3,i)_{(v_0,v_3)}$-path for any $i \in C \backslash \{3\}$ and thus G contains no $(3,6)_{(v_0,v_1)}$-cycle. (See (a), (b) in Fig. 6.)

If $4 \in C(v_0)$, then let $\{vv_1, v_2v_3\} \to 2$, $(v_1v_0, vv_0) \to (1,3)$. If $4 \notin C(v_3)$, then let $(vv_2, vv_3, v_1v_2) \to (1,4,6)$. Otherwise, $4 \in C(v_3)$ and $5 \in C(v_3) \cap C(v_0)$ similarly. It follows that $C(v_3) = C(v_0) = \{4,5\}$. Then, let $(vv_3, v_1v_2, vv_2) \to (1,3,6)$. Since $c(v_0v_3) = 3$ in H, it is obvious that $3 \notin C(v_3)$ and thus G contains no $(3,2)_{(v_0,v)}$-cycle. (See (a), (c) in Fig. 6.)

Case 3 $v_1v_2, v_2v_3 \in E(G)$ and $v_1v_0, v_3v_0 \notin E(G)$, as shown in Fig. 7.

If $v_1v_3, v_2v_0 \in E(G)$, then $v_1v_3, v_3v_2, v_2v_0 \in E(G)$ and the proof can be reduced to Case 2. Otherwise, $v_1v_3 \notin E(G)$ or $v_2v_0 \notin E(G)$.

Case 3.1 $v_1v_3 \in E(G)$ and $v_2v_0 \notin E(G)$.



(a) H (b) (c)

Fig. 6: Case 2.2.2.



Case 3.1 Case 3.2 Case 3.3

Fig. 7: The configuration in Case 3.

Let v_5 be the neighbor of v_1 different from v, v_2, v_3. Let v_6 be the neighbor of v_2 different from v, v_1, v_3. Let v_7 be the neighbor of v_3 different from v, v_1, v_2. By Cases 1 and 2, we can assume that v_0, v_5, v_6, v_7 are pairwise distinct.

Let $H = G - \{v, v_1, v_2, v_3\} + \{u, uv_0, uv_5, uv_6, uv_7\}$ and assume that $(uv_5, uv_6, uv_7, uv_0)_c = (1, 2, 3, 4)$. Then, let $\{v_1v_5, v_2v_3\} \to 1$, $\{vv_3, v_2v_6\} \to 2$, $\{vv_1, v_3v_7\} \to 3$, $\{vv_2, v_1v_3\} \to 5$, and $(vv_0, v_1v_2) \to (4, 6)$.

Case 3.2 $v_1v_3 \notin E(G)$ and $v_2v_0 \in E(G)$.

Let $H = G - \{v, v_2\} + \{v_1v_0, v_0v_3, v_3v_1\}$ and assume that $(v_1v_0, v_0v_3, v_3v_1)_c = (1, 2, 3)$, $4 \in C \backslash (C(v_1) \cup \{1, 2, 3\})$. Then, first let $\{vv_3, v_1v_2\} \to 3$, $(vv_0, vv_1) \to (1, 4)$. Next, if $\{5, 6\} \backslash C(v_3) \neq \emptyset$ and $5 \notin C(v_3)$, then let $(v_2v_0, v_2v_3, vv_2) \to (2, 5, 6)$; otherwise, $C(v_3) = \{5, 6\}$ and let $v_2v_3 \to 1$. If $\{5, 6\} \backslash C(v_0) \neq \emptyset$, then let $v_2v_0 \to 2$ and $vv_2 \to \beta \in \{5, 6\} \backslash C(v_0)$; otherwise, $C(v_0) = \{5, 6\}$ and let $(vv_2, v_2v_0) \to (2, 4)$.

Case 3.3 $v_1v_3 \notin E(G)$ and $v_2v_0 \notin E(G)$.

Let v_5 be the neighbor of v_2 different from v, v_1, v_3. Let v_6, v_7 be the other two neighbors of v_1 different from v, v_2. Note that v_0, v_5, v_6, v_7 are pairwise distinct by the previous argument. Let $H = G - \{v, v_2\} + \{v_1v_0, v_1v_5\}$ and assume that $(v_1v_5,\ v_1v_0,\ v_1v_6,\ v_1v_7)_c = (1, 2, 3, 4)$. Then, first let $(v_2v_5, vv_0) \to (1, 2)$.

Case 3.3.1 $C(v_3) = \{1, 2\}$.

Assume that $6 \notin C(v_5)$. Then, first let $(v_1v_2, vv_2, v_2v_3) \to (2, 4, 6)$. Next, if $5 \notin C(v_0)$, then let $(vv_1, vv_3) \to (1, 5)$; if $3 \notin C(v_0)$, then let $(vv_1, vv_3) \to (5, 3)$; otherwise, $C(v_0) = \{2, 5, 3, 4\}$ and let $(vv_1, vv_3, vv_0) \to (1, 5, 6)$.

Assume that $\{5, 6\} \subseteq C(v_5)$ and $\{5, 6\} \subseteq C(v_0)$ similarly. If $3, 4 \notin C(v_0) \cup C(v_5)$, then let $\{vv_1, v_2v_3\} \to 5$ and $(v_1v_2, v_2v_5, vv_3, vv_2) \to (1, 3, 4, 6)$; otherwise, $C(v_0) = \{2, 5, 6, 4\}$. Then, let $(vv_1, vv_2, vv_0, vv_3, v_1v_2) \to (1, 2, 3, 5, 6)$ and $v_2v_3 \to \gamma \in \{3, 4\} \backslash C(v_5)$.

Case 3.3.2 $C(v_3) = \{1, 3\}$.

If $6 \notin C(v_5)$, then let $\{vv_1, v_2v_3\} \to 6$ and $(v_1v_2, vv_3, vv_2) \to (2, 4, 5)$. If G contains no $(1, 4)_{(v_3,v_5)}$-path, then let $(vv_1, vv_2, v_2v_3, vv_3, v_1v_2) \to (1, 3, 4, 5, 6)$. Otherwise, G contains a $(1, 4)_{(v_3,v_5)}$-path and hence $C(v_5) = \{1, 5, 6, 4\}$. We let $(vv_1, v_2v_3, vv_3, v_1v_2, vv_2) \to (1, 2, 4, 5, 6)$.

Case 3.3.3 $C(v_3) = \{1, 5\}$.

Let $vv_1 \to 6$. If $6 \notin C(v_5)$, we further let $(v_1v_2, vv_3, vv_2,\ v_2v_3) \to (2, 3, 5, 6)$; If $3 \notin C(v_5)$, let $(v_1v_2, v_2v_3, vv_3, vv_2) \to (2, 3, 4, 5)$. Otherwise, $C(v_5) = \{1, 6, 3, 4\}$, let $(v_2v_3, vv_3, vv_2, v_1v_2) \to (2, 3, 4, 5)$.

Case 3.3.4 $C(v_3) \notin \{\{1, 2\}, \{1, 3\}, \{1, 5\}\}$.

If $C(v_3) = \{5, 6\}$, let $(vv_1, v_2v_3, vv_3, vv_2, v_1v_2) \to (1, 2, 3, 4, 5)$. If $C(v_3) = \{3, 5\}$, let $\{vv_1, v_2v_3\} \to 6$ and $(vv_3, vv_2, v_1v_2) \to (1, 4, 5)$. If $C(v_3) = \{3, 4\}$, let $\{vv_1, v_2v_3\} \to 5$ and $(vv_3, v_1v_2, vv_2) \to (1, 2, 6)$.

Case 4 $v_1v_2, v_3v_0 \in E(G)$ and $v_0v_1, v_2v_3, v_1v_3, v_2v_0 \notin E(G)$.

Let $V_i = \{v_{i1}, v_{i2}\}$ be the set of the other neighbors of v_i for $i \in \{0, 1, 2, 3\}$. By Cases 1 to 3, $V_1 \cap V_2 = \emptyset$ and $V_3 \cap V_0 = \emptyset$. Let $H = G - \{v, v_1, v_2, v_3, v_0\} + \{u, w, uv_{11}, uv_{12}, uv_{21}, uv_{22}, wv_{31}, wv_{32}, wv_{01}, wv_{02}\}$, where u and w are new vertices added. Assume that $(uv_{11},\ uv_{12},\ uv_{21},\ uv_{22})_c = (1,\ 2,\ 3,\ 4)$, $\{c(wv_{01}), c(wv_{02})\} = \{a, b\}$, $\{c(wv_{31}), c(wv_{32})\} = \{c, d\}$, where $\{a, b\} \cap \{c, d\} = \emptyset$, see Fig. 8. Then, first let $v_iv_{ij} \to c(uv_{ij})$ for $i \in \{1, 2\}$, and $vv_{ij} \to c(wv_{ij})$ for $i \in \{3, 4\}$, $j \in \{1, 2\}$. Next, by symmetry, we need to consider the following.

Case 4.1 $\{a, b\} = \{1, 3\}$ and $\{c, d\} = \{2, 4\}$.



Case 4

H

Fig. 8: The configuration in Case 4.

If $5 \notin C(v_{11})$, let $\{v_1v_2, vv_0\} \to 5$ and $(vv_2, vv_3, vv_1, v_3v_0) \to (1,3,4,6)$. If $4 \notin C(v_{11})$, let $\{v_1v_2, vv_0\} \to 5$ and $(vv_3, vv_2, vv_1, v_3v_0) \to (1,2,4,6)$. Otherwise, $C(v_{11}) = \{1,4,5,6\}$ and $C(v_{12}) = \{2,3,5,6\}$. Let $\{v_1v_2, vv_3\} \to 1$, $\{vv_1, v_3v_0\} \to 5$, and $(vv_2, v_1v_{11}, vv_0) \to (2,3,6)$.

Case 4.2 $\{a,b\} = \{1,2\}$ and $\{c,d\} = \{3,4\}$.

If G contains no $(i,j)_{(v_2,v_0)}$-path for some $i \in \{1,2\}$ and $j \in \{3,4\}$, then let $\{v_1v_2, vv_3\} \to 5$, $\{vv_1, v_3v_0\} \to 6$, and $(vv_2, vv_0) \to (i,j)$. Otherwise, G contains a $(i,j)_{(v_2,v_0)}$-path for any $i \in \{1,2\}$ and $j \in \{3,4\}$ and $\{1,2\} \subseteq C(v_{21}) \cap C(v_{22})$. It follows that $\{5,6\} \backslash C(v_{21}) \neq \emptyset$ and assume that $5 \notin C(v_{21})$. First, let $\{vv_2, v_3v_0\} \to 5$, $\{v_1v_2, vv_3\} \to 6$, and $(vv_0, vv_1) \to (3,4)$. Next, if $6 \notin C(v_{22})$, then we are done; otherwise, $C(v_{22}) = \{4,1,2,6\}$ and let $\{vv_1, v_2v_{22}\} \to 5$, $vv_2 \to 4$.

Case 4.3 All other cases.

If $\{a,b\} = \{1,2\}$ and $\{c,d\} = \{5,6\}$, let $\{vv_1, v_3v_0\} \to 3$ and $(vv_2,\ vv_3,\ v_1v_2,\ vv_0) \to (1,\ 4,\ 5,\ 6)$. If $\{a,b\} = \{1,3\}$ and $\{c,d\} = \{5,6\}$, let $(vv_2,\ vv_0,\ vv_3,\ v_3v_0, v_1v_2, vv_1) \to (1,2,3,4,5,6)$. If $\{a,b\} = \{1,5\}$ and $\{c,d\} = \{2,6\}$, let $(vv_2,\ vv_0,\ v_3v_0,\ vv_3, v_1v_2, vv_1) \to (1,2,3,4,5,6)$. If $\{a,b\} = \{1,5\}$ and $\{c,d\} = \{3,6\}$, let $\{v_1v_2, vv_3\} \to 5$ and $(vv_2, vv_0, v_3v_0, vv_1) \to (1,2,4,6)$. If $\{a,b\} = \{1,2\}$ and $\{c,d\} = \{3,5\}$, let $\{v_1v_2, v_3v_0\} \to 6$ and $(vv_2, vv_1, vv_3, vv_0) \to (1,3,4,5)$. If $\{a,b\} = \{1,3\}$ and $\{c,d\} = \{2,5\}$, let $\{v_1v_2, v_3v_0\} \to 6$ and $(vv_2, vv_0, vv_3, vv_1) \to (1,2,4,5)$.

Case 5 $v_1v_2 \in E(G)$ and $v_2v_3, v_3v_0, v_0v_1, v_1v_3, v_2v_0 \notin E(G)$.

Let V_1, V_2 be defined similarly as in Case 4. By Cases 1 to 4 , $V_1 \cap V_2 = \emptyset$. Let $H = G - \{v, v_1, v_2\} + \{u, uv_{11}, uv_{12}, uv_{21}, uv_{22}, v_3v_0\}$ and assume $(uv_{11}, uv_{12}, uv_{21}, uv_{22})_c = (1,2,3,4)$, as shown in Fig. 9. Then, first let $v_iv_{ij} \to c(uv_{ij})$ for $i,j \in \{1,2\}$. Next, since $c(v_3v_0) = 1$ or $c(v_3v_0) = 5$, we need to consider the following.

Case 5.1 $c(v_3v_0) = 1$.



Case 5 H

Fig. 9: The configuration in Case 5.

Case 5.1.1 $2 \notin C(v_3)$.

If G contains neither $(2,3)_{(v_1,v_3)}$-path nor $(1,3)_{(v_1,v_0)}$-path, let $(vv_0, vv_3, vv_1, v_1v_2, vv_2) \to (1,2,3,5,6)$. Otherwise, G contains either a $(2,3)_{(v_1,v_3)}$-path or a $(1,3)_{(v_1,v_0)}$-path. If $3 \in C(v_3)\backslash C(v_0)$, then let $(vv_2, vv_3, vv_0) \to (1,2,3)$ and $vv_1 \to x_1 \in \{4,5,6\}\backslash C(v_3)$, $v_1v_2 \to y_1 \in \{5,6\}\backslash\{x_1\}$. If $3 \in C(v_0)\backslash C(v_3)$, then let $(vv_0, vv_2, vv_3) \to (1,2,3)$ and $vv_1 \to x_2 \in \{4,5,6\}\backslash C(v_0)$, $v_1v_2 \to y_2 \in \{5,6\}\backslash\{x_2\}$. Otherwise, $3 \in C(v_3) \cap C(v_0)$, $4 \in C(v_3) \cap C(v_0)$ similarly and then $5 \notin C(v_3)$. First, let $\{v_1v_2, vv_3\} \to 5$ and $(vv_0, vv_2, vv_1) \to (1,2,6)$. Next, if G contains no $(1,6)_{(v_1,v_0)}$-path, then we are done; otherwise, G contains a $(1,6)_{(v_1,v_0)}$-path, which cannot pass through v_3 and $C(v_0) = \{3,4,6\}$. We switch the colors of vv_3 and vv_0. Otherwise, assume that $2 \in C(v_3)$ and $2 \in C(v_0)$ similarly.

Case 5.1.2 $3 \notin C(v_3)$.

First, assume that G contains no $(1,4)_{(v_1,v_0)}$-path and let $(vv_0, vv_3, vv_1) \to (1,3,4)$. If G contains no $(3,i)_{(v_2,v_3)}$-path for some $i \in \{2,5,6\}$, then let $vv_2 \to i$ and $v_1v_2 \to \alpha \in \{5,6\}\backslash\{i\}$. Otherwise, G contains a $(3,i)_{(v_2,v_3)}$-path for any $i \in \{2,5,6\}$ and $C(v_{21}) = \{3,2,5,6\}$, $C(v_3) = \{2,5,6\}$. If $\{5,6\}\backslash C(v_0) \neq \emptyset$ and $5 \notin C(v_0)$, then let $(v_2v_{21}, v_1v_2, vv_2) \to (1,3,5)$; otherwise, $\{5,6\} \subseteq C(v_0)$, $C(v_0) = \{1,2,5,6\}$ and let $(vv_2, vv_0, v_1v_2, vv_1) \to (1,4,5,6)$. Next, assume that G contains a $(1,4)_{(v_1,v_0)}$-path and $4 \in C(v_0)$, $5 \notin C(v_0)$. Then, let $(vv_3, vv_2, vv_1, vv_0, v_1v_2) \to (1,2,4,5,6)$. Otherwise, $3 \in C(v_3)$ and $3 \in C(v_0)$.

Case 5.1.3 $C(v_3) = C(v_0) = \{2,3,4\}$.

First, let $\{v_1v_2, vv_0\} \to 5$ and $(vv_3, vv_2, vv_1) \to (1,2,6)$. Next, if G contains no $(2,5)_{(v_1,v_0)}$-path, then we are done; otherwise, G contains a $(2,5)_{(v_1,v_0)}$-path, we switch the colors of vv_3 and vv_0.

Case 5.2 $c(v_3v_0) = 5$.

By symmetry, we may assume that $1 \notin C(v_3)$. If G contains no $(1,i)_{(v_1,v_3)}$-path for some $i \in \{3,4\}$, then let $(vv_3, vv_2, vv_1, vv_0, v_1v_2) \to (1,2,i,5,6)$. Otherwise, G contains a $(1,i)_{(v_1,v_3)}$-path for any $i \in \{3,4\}$ and $\{3,4\} \subseteq C(v_{11}) \cap C(v_3)$. If $1 \notin C(v_0)$, then let $(vv_0, vv_2, vv_1, vv_3, v_1v_2) \to (1,2,3,5,6)$. Otherwise, $1 \in C(v_0)$.

Case 5.2.1 G contains no $(2,5)_{(v_1,v_0)}$-path.

If G contains no $(1,6)_{(v_1,v_3)}$-path, then let $\{v_1v_2, vv_0\} \to 5$ and $(vv_3, vv_2, vv_1) \to (1,2,6)$. Otherwise, G contains a $(1,6)_{(v_1,v_3)}$-path and $6 \in C(v_{11}) \cap C(v_3)$ and $C(v_{11}) = \{1,3,4,6\}$, $C(v_3) = \{3,4,6\}$. If G contains no $(2,i)_{(v_1,v_3)}$-path for some $i \in \{3,4,6\}$, then let $(vv_2, vv_3, vv_4, vv_1) \to (1,2,5,i)$ and $v_1v_2 \to x_4 \in \{5,6\} \backslash \{i\}$. Otherwise, G contains a $(2,i)_{(v_1,v_3)}$-path for any $i \in \{3,4,6\}$ and $C(v_{12}) = \{2,3,4,6\}$. If $2 \notin C(v_0)$, then let $(vv_2, vv_0, vv_1, vv_3, v_1v_2) \to (1,2,3,5,6)$. If $6 \notin C(v_0)$, then let $\{v_1v_2, vv_0\} \to 6$ and $(vv_3, vv_2, vv_1) \to (1,2,5)$. Otherwise, $C(v_0) = \{1,2,6\}$ and $4 \notin C(v_0)$. Then, let $\{v_1v_2, vv_3\} \to 1$, $\{vv_0, v_1v_{11}\} \to 5$, and $(vv_1, vv_2) \to (4,6)$.

Case 5.2.2 G contains a $(2,5)_{(v_1,v_0)}$-path, and $5 \in C(v_{12})$, $2 \in C(v_0)$.

Then, $\{3,4\} \backslash C(v_0) \neq \emptyset$ and assume that $3 \notin C(v_0)$. If $5 \notin C(v_{11})$, then let $\{vv_0, v_1v_{11}\} \to 5$ and $(vv_3, vv_2, vv_1, v_1v_2) \to (1,2,3,6)$. Otherwise, $C(v_{11}) = \{1,3,4,5\}$. Then, let $\{v_1v_2, vv_3\} \to 1$ and $(vv_0, vv_1, vv_2, v_1v_{11}) \to (3,4,5,6)$. □

References

[1] N. Alon, B. Sudakov, A. Zaks, Acyclic edge colorings of graphs, J. Graph Theory 37 (2001) 157-167.

[2] N. Alon, C. McDiarmid, B. Reed, Acyclic coloring of graphs, Random Structures Algorithms 2 (1991) 277-288.

[3] M. Basavaraju, L.S. Chandran, Acyclic edge coloring of subcubic graphs, Discrete Math. 308 (2008) 6650-6653.

[4] M. Basavaraju, L.S. Chandran, Acyclic edge coloring of graphs with maximum with degree 4, J. Graph Theory 61 (2009) 192-209.

[5] M. Basavaraju, L.S. Chandran, Acyclic edge coloring of 2-degenerate graphs, J. Graph Theory 69 (2012) 1-27.

[6] M. Basavaraju M, L.S. Chandran, N. Cohen, F. Havet, T. Müller, Acyclic edge-coloring of planar graphs, SIAM J. Discrete Math. 25 (2011) 463-478.

[7] N. Cohen N, F. Havet, T. Müller, Acyclic edge-colouring of planar graphs, Extended abstract, Electron. Notes Discrete Math. 34 (2009) 417-421.

[8] W. Dong, B. Xu, Some results on acyclic edge coloring of plane graphs, Inform. Process. Lett. 110 (2010) 887-892.

[9] L. Esperet, A. Parreau, Acyclic edge-coloring using entropy compression, arXiv: 1206.1535v1 [math.CO], 7 Jun 2012.

[10] J. Fiamčik, The acyclic chromatic class of a graph, Math Slovaca 28 (1978) 139-145 (in Russian).

[11] A. Fiedorowicz, M. Haluszczak, N. Narayanan, About acyclic edge colourings of planar graphs, Inform. Process. Lett. 108 (2008) 412-417.

[12] J. Hou, J. Wu, G. Liu, B. Liu, Acyclic edge colorings of planar graphs and series-parappel graphs, Sci. China Ser. A 52 (2009) 605-616.

[13] J. Hou, J. Wu, G. Liu, B. Liu B, Acyclic edge chromatic number of outerplanar graphs, J. Graph Theory 64 (2010) 22-36.

[14] R. Muthu, N. Narayanan, C. R. Subramanian, Acyclic edge colouring of outerplanar graphs, Lecture Notes Comput. Sci. 4508 (2007) 144-152.

[15] M. Molloy, B. Reed, Further algorithmic aspects of Lovász local lemma, in Proceedings of the 30th Annual ACM Symposium on Theory of Computing, Dallas, Texas, United States, 1998, pp. 524-529.

[16] S. Ndreca, A. Procacci, B. Scoppola, Improved bounds on coloring of graphs, European J. Combin. 33 (2012) 592-609.

[17] J. Něsetřil, N.C. Wormald, The acyclic edge chromatic number of a random d-regular graph is $d+1$, J. Graph Theory 49 (2005) 69-74.

[18] S. Skulrattanakulchai, Acyclic colorings of subcubic graphs, Inform. Process. Lett. 92 (2004) 161-167.

[19] Q. Shu, W. Wang, Acyclic chromatic indices of planar graphs with girth at least five, J. Comb. Optim. 23 (2012) 140-157.

[20] Q. Shu, W. Wang, Y. Wang, Acyclic chromatic indices of planar graphs with girth at least four, J. Graph Theory, DOI: 10.1002/jgt.21683.

[21] Q. Shu, W. Wang, Y. Wang, Acyclic edge coloring of planar graphs without 5-cycles, Discrete Appl. Math. 160 (2012). 1211-1223.

[22] W. Wang, Q. Shu, Acyclic chromatic indices of K_4-minor free graphs, Sci. China Ser. A 41 (2011) 733-744.

[23] W. Wang, Q. Shu, W. Kan, P. Wang, Acyclic chromatic indices of planar graphs with large girth, Discrete Appl. Math. 159 (2011) 1239-1253.

[24] W. Wang, Q. Shu, Y. Wang, Acyclic edge coloring of planar graphs without 4-cycles, J. Comb. Optim. DOI: 10.1007/s10878-012-9474-y.

[25] W. Wang, Q. Shu, Y. Wang, A new upper bound on the acyclic chromatic indices of planar graphs, European J. Combin., In press.

[26] V. Vizing, On an estimate of the chromatic index of a p-graph, Diskret Analiz 3 (1964) 25-30.

[27] D. Yu, J. Hou, G. Liu, B. Liu, L. Xu, Acyclic edge coloring of planar graphs with large girth, Theoret. Comput. Sci. 410 (2010) 5196-5200.